

A DETAILED STUDY OF VARIOUS GRADES OF ANAEMIA IN PATIENTS WITH HOOKWORM INFECTION DIAGNOSED BY DOING UPPER GASTRO INTESTINAL ENDOSCOPY

GOVINDARAJALU GANESAN*

**Associate professor, Dept. of General surgery, Indira Gandhi Medical College and Research Institute,
Puducherry.-605009*

ABSTRACT:

Objective: Anaemia is commonly reported to occur in hookworm infection. Hence a detailed study was done to know about the various grades of anaemia in patients with hookworm infection diagnosed by doing upper gastro intestinal endoscopy.

Methods: A study of 1307 patients who had undergone upper gastro-intestinal endoscopy for a period of 5 years and one month from May 2009 to May 2014 was carried out to know about the various grades of anaemia in patients with hookworm infection diagnosed by doing upper gastro intestinal endoscopy.

Results: Out of these 1307 patients, 14 patients found to have hookworms in duodenum while doing upper gastro-intestinal endoscopy were taken into consideration for our study. Out of these 14 patients with hookworms in duodenum, 5 patients did not have anaemia, 6 patients had mild anaemia, 1 patient had moderate anaemia and 2 patients were found to have severe anaemia. Hence 11 out of 14 patients with hookworms in duodenum had absence of anaemia and mild anaemia. Hence majority of the patients with hookworms in duodenum were in the early stage of hookworm infection in our study since 11 out of 14 patients had absence of anaemia and only mild anaemia. Only 3 out of 14 patients with hookworms in duodenum had moderate anaemia and severe anaemia. Hence only few patients with hookworms in duodenum were in the late stage of hookworm infection in our study since only 3 out of 14 patients had moderate anaemia and severe anaemia.

Conclusion: Hence majority of the patients with hookworms in duodenum were in the early stage of hookworm infection in our study. Only few patients with hookworms in duodenum were in the late stage of hookworm infection in our study.

Keywords: various grades of anaemia, hookworms in duodenum, upper gastro intestinal endoscopy

INTRODUCTION:

Anaemia is commonly reported to occur in hookworm infection (1 to 16). Hence a detailed study was done to know about the various grades of anaemia in patients with hookworm infection diagnosed by doing upper gastro intestinal endoscopy.

MATERIALS AND METHODS:

This study was conducted in the department of general surgery, Aarupadai Veedu Medical College And Hospital, Puducherry. A study of 1307 patients who had undergone upper gastro-intestinal endoscopy for a period of 5 years and one month from May 2009 to May 2014 was carried out. In each of these 1307 patients, the first and second part of duodenum was carefully examined to find out the presence of hookworms. In all the patients found to have hookworms in duodenum, investigations were done to know about the presence or absence of anaemia and to know about the various grades of anaemia in patients with hookworm infection diagnosed while doing upper gastro intestinal endoscopy. Anaemia is defined as hemoglobin <12g/dl or 12g% in women and hemoglobin or <13g/dl or 13g% in men. Mild anaemia is taken as hemoglobin 10 to 12g/dl or g%, moderate anaemia is taken as haemoglobin 7 to 10g/dl or g% and severe anaemia is taken as hemoglobin <7g/dl or g%. The results were found as given below.

RESULTS:

1. Out of these 1307 patients, 14 patients found to have hookworms in duodenum while doing upper gastro-intestinal endoscopy were taken into consideration for our study.
2. Out of these 14 patients with hookworms in duodenum taken into consideration for our study.
 - a. 5 patients did not have anaemia [hemoglobin >13g/dl in men and hemoglobin >12g/dl in women].
 - b. 6 patients had mild anaemia [haemoglobin 10 to 12g/dl]
 - c. 1 patient had moderate anaemia [haemoglobin 7 to 10g/dl]
 - d. 2 patients were found to have severe anaemia [hemoglobin <7g/dl].
3. 11 out of 14 patients with hookworms in duodenum had absence of anaemia and only mild anaemia.
4. Only 3 out of 14 patients with hookworms in duodenum had moderate anaemia and severe anaemia.

DISCUSSION:

Absence of anaemia

Out of the 14 patients with hookworms in duodenum taken into consideration for our study, 5 patients did not have anaemia [haemoglobin >13g/dl in men and haemoglobin >12g/dl in women].

Single hookworm in duodenum seen in one of these patients with absence of anaemia [haemoglobin 17.6g%] is shown in Fig 1.

The hookworm in duodenum is identified by its bent head which looks like a hook (Fig 1) and by its S-shaped appearance (13) (Fig 1).

Mild anaemia

Out of the 14 patients with hookworms in duodenum taken into consideration for our study, 6 patients had mild anaemia [haemoglobin 10 to 12g/dl].

Single hookworm in duodenum seen in one of these patients with mild anaemia [haemoglobin 10g/dl or g %] is shown in Fig 2.

The hookworm in duodenum is identified by its bent head which looks like a hook (Fig 1) and by its S-shaped appearance (13) (Fig 1).

Moderate anaemia

Out of the 14 patients with hookworms in duodenum taken into consideration for our study, only 1 patient had moderate anaemia [haemoglobin 7 to 10g/dl].

Severe anaemia

Out of the 14 patients with hookworms in duodenum taken into consideration for our study, only 2 patients were found to have severe anaemia [haemoglobin <7g/dl].

Multiple hookworms in duodenum seen in one of these 2 patients with severe anaemia [haemoglobin 3.2 g/dl or g %] is shown in fig 3.

Hence 11 out of 14 patients (79%) with hookworms in duodenum had absence of anaemia and mild anaemia.

But only 3 out of 14 patients (21%) with hookworms in duodenum had moderate anaemia and severe anaemia.




Fig1: Single hookworm in duodenum with its bent head and S-shaped appear ancient patient Without anaemia [haemoglobin 17.6g%]




Fig2:Single hookworm in duodenum with its bent head like a hook and S-shaped appearance
In a patient with mild anaemia [haemoglobin 10 g%]



Fig 3: Multiple hookworms in duodenum in a patient with severe anaemia [haemoglobin 3.2 g%].

CONCLUSION:

1. Hence majority of the patients with hookworms in duodenum were in the early stage of hookworm infection in our study since 11 out of 14 patients with hookworms in duodenum had absence of anaemia and only mild anaemia.
2. Hence only few patients with hookworms in duodenum were in the late stage of hookworm infection in our study since only 3 out of 14 patients with hookworms in duodenum had moderate anaemia and severe anaemia.

ACKNOWLEDGEMENT:

The author acknowledges the immense help received from the scholars whose articles are cited and included in references of this manuscript. The author is also grateful to authors / editors / publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed. The author is extremely grateful to IJCRR editorial board members and IJCRR team of reviewers who have helped to bring quality to this manuscript.

REFERENCES

1. Hyun HJ, Kim EM, Park SY, Jung JO, Chai JY, Hong ST .A case of severe anemia by *Necator americanus* infection in Korea. J Korean Med Sci. 2010 Dec;25(12):1802-4.
2. Wu KL, Chuah SK, Hsu CC, Chiu KW, Chiu YC, Changchien CS. Endoscopic Diagnosis of Hookworm Disease of the Duodenum: A Case Report. J Intern Med Taiwan 2002;13:27-30.
3. Kuo YC, Chang CW, Chen CJ, Wang TE, Chang WH, Shih SC . Endoscopic Diagnosis of Hookworm Infection That Caused Anemia in an Elderly Person. International Journal of Gerontology. 2010 ; 4(4) : 199-201
4. Nakagawa Y, Nagai T, Okawara H, Nakashima H, Tasaki T, Soma W, et al. Comparison of magnified endoscopic images of *Ancylostoma duodenale* (hookworm) and *Anisakis simplex*. Endoscopy 2009;41(Suppl. 2):E189
5. Basset D, Rullier P, Segalas F, Sasso M. Hookworm discovered in a patient presenting with severe iron-deficiency anemia. Med Trop (Mars). 2010 Apr;70(2):203-4
6. Lee T.-H., Yang J.-c., Lin J.-T., Lu S.-C. and Wang T.-H. Hookworm Infection Diagnosed by Upper Gastrointestinal Endoscopy: —Report of Two Cases with Review of the Literature—. Digestive Endoscopy, 1994 6(1): 66–72
7. Anjum Saeed, Huma Arshad Cheema, Arshad Alvi, Hassan Suleman. Hookworm infestation in children presenting with melena -case series. Pak J Med Res Oct - Dec 2008;47(4) :98-100

8. A Rodríguez, E Pozo, R Fernández, J Amo, T Nozal. Hookworm disease as a cause of iron deficiency anemia in the prison population Rev EspSanidPenit 2013; 15: 63-65
9. Li ZS¹, Liao Z, Ye P, Wu RP Dancing hookworm in the small bowel detected by capsule endoscopy: a synthesized video. Endoscopy. 2007 Feb;39Suppl 1:E97. Epub 2007 Apr 18.
10. Kalli T¹, Karamanolis G, Triantafyllou K Hookworm infection detected by capsule endoscopy in a young man with iron deficiency. ClinGastroenterolHepatol. 2011 Apr;9(4):e33
- 11.Chen JM¹, Zhang XM, Wang LJ, Chen Y, Du Q, Cai JT. Overt gastrointestinal bleeding because of hookworm infection. Asian Pac J Trop Med. 2012 Apr;5(4):331-2.
12. Kato T, Kamoi R, Iida M, KiharaT.Endoscopic diagnosis of hookworm disease of the duodenum J ClinGastroenterol. 1997 Mar;24(2):100-102
13. Cedrón-Cheng H, Ortiz C Hookworm Infestation Diagnosed by Capsule Endoscopy. J Gastroint Dig Syst2011S1:003. doi: 10.4172/2161-069X.S1-003
14. Yan SL, Chu YC. Hookworm infestation of the small intestine Endoscopy 2007; 39: E162±163
15. Chao CC¹, Ray ML. Education and imaging. Gastrointestinal: Hookworm diagnosed by capsule endoscopy. J GastroenterolHepatol. 2006 Nov;21(11):1754.
16. Christodoulou, D. K., Sigounas, D. E., Katsanos, K. H., Dimos, G., &Tsianos, E. V. (). Small bowel parasitosis as cause of obscure gastrointestinal bleeding diagnosed by capsule endoscopy.World journal of gastrointestinal endoscopy, 2(11), 2010: 369.